

**75061**  
Mare Soil  
186.5 grams




Figure 1: Map of Apollo 17 showing location of soil samples and coarse-fines (Meyer 1973). S73-24071




Figure 2: Map of station 5, Apollo 17.

**Introduction**

75060 was collected from the top surface of a flat boulder of basalt on the rim of Camelot Crater. It is made up of comminuted pieces of that boulder mixed with some soil that was undoubtedly kicked up by meteorite impacts in the adjacent soil. 75075 is a large piece of the boulder and 75080 is a sample of nearby soil (also very basaltic).

**Petrography**

The maturity of 75061 is  $I_s/FeO = 33$  and the average grain size is 112 microns (Morris 1978, Graf 1993). Heiken and McKay (1974) reported on 24% agglutinate. Meyer (1973) found that most all of the 4 – 10 mm coarse fines were mare basalt.

**Chemistry**

The FeO content of 75061 is very high (figure 3), and the composition is similar to the basaltic boulder (table).

Muller (1974) determined 42 ppm nitrogen in 75061. Petrowski et al. (1974) determined 94 ppm carbon, 49 ppm nitrogen and 43 ppm hydrogen




Figure 3: Composition of 75061 compared with other Apollo soils.




Figure 4: Normalized rare-earth-element diagram for 75061 compared with highland soil 72221.





Figure 5: Carbon content and maturity index for 75061.

$^{48}\text{V} = 26 \text{ dpm/kg}$ ,  $^{54}\text{Mn} = 286 \text{ dpm/kg}$ ,  $^{7}\text{Be} = 350 \text{ dpm/kg}$  and  $^{56}\text{Co} = 548 \text{ dpm/kg}$ .

### Cosmogenic isotopes and exposure ages

Keith et al. (1974) determined the cosmic-ray-induced activity of  $^{22}\text{Na} = 187 \text{ dpm/kg}$ ,  $^{26}\text{Al} = 180 \text{ dpm/kg}$ ,  $^{46}\text{Sc} = 86 \text{ dpm/kg}$ ,  $^{48}\text{V} = 47 \text{ dpm/kg}$ ,  $^{54}\text{Mn} = 200 \text{ dpm/kg}$  and  $^{56}\text{Co} = 490 \text{ dpm/kg}$ . Rancitelli et al. (1974) determined the cosmic-ray-induced activity of  $^{22}\text{Na} = 171 \text{ dpm/kg}$ ,  $^{26}\text{Al} = 174 \text{ dpm/kg}$ ,  $^{46}\text{Sc} = 112 \text{ dpm/kg}$ ,



average grain size = 112 microns

### Modal content of soil 75061 (90-150 micron).

From Heiken and McKay 1974.

	75061
Agglutinates	24 %
Basalt	26.6
Breccia	4.9
Anorthosite	
Norite	
Gabbro	
Plagioclase	4.6
Pyroxene	29.6
Olivine	0.3
Ilmenite	5.3
Orange glass	1
Glass other	3.2





Figure 6: Grain size distribution of 75061 (Graf 1993, data from McKay).

**Table 1. Chemical composition of 75061**

reference weight	LSPET73 Rhodes74	Wiesman76 Rhodes74	Korotev92		Rose74	Keith74	Rancitelli74	Brunfelt74 unpublished	
SiO <sub>2</sub> %	39.32 (a)				39.7 (d)				
TiO <sub>2</sub>	10.31 (a)	11 (b)			10.46 (d)			9.5 (c)	
Al <sub>2</sub> O <sub>3</sub>	10.42 (a)				10.6 (d)			10.5 (c)	
FeO	18.19 (a)		18	17.9 (c)	17.86 (d)			16.8 (c)	
MnO	0.25 (a)				0.24 (d)			0.24 (c)	
MgO	9.53 (a)				9.65 (d)			7.8 (c)	
CaO	10.72 (a)				10.72 (d)			8.96 (c)	
Na <sub>2</sub> O	0.33 (a)	0.35	0.344	0.36 (c)	0.37 (d)			0.39 (c)	
K <sub>2</sub> O	0.08 (a)	0.075 (b)			0.08 (d)	0.079	0.072 (e)	0.084 (c)	
P <sub>2</sub> O <sub>5</sub>	0.06 (a)								
S %	0.13 (a)								
sum									
Sc ppm			75	70 (c)	78 (d)			65 (c)	
V					95 (d)			137 (c)	
Cr	3284 (a)	3090 (b)	3480	3280 (c)	3284 (d)			2840 (c)	
Co			26.3	27.8 (c)	36 (d)			25 (c)	
Ni	115 (a)		100	100 (c)	154 (d)				
Cu					27 (d)			6.9 (c)	
Zn	25 (a)				12 (d)			30 (c)	
Ga					4.3 (d)			4.1 (c)	
Ge ppb									
As									
Se									
Rb	1.6 (a)	1.11 (b)			1 (d)			1.3 (c)	
Sr	166 (a)	1.66 (b)	130	210 (c)	157 (d)			173 (c)	
Y	83 (a)				91 (d)				
Zr	237 (a)		250	330 (c)	340 (d)				
Nb	21 (a)				20 (d)				
Mo									
Ru									
Rh									
Pd ppb									
Ag ppb									
Cd ppb									
In ppb									
Sn ppb									
Sb ppb									
Te ppb									
Cs ppm								0.075 (c)	
Ba		89.5 (b)	80	105 (c)	135 (d)			76 (c)	
La		7.07 (b)	6.4	7.34 (c)				5.7 (c)	
Ce		23.6 (b)	21	21.3 (c)					
Pr									
Nd		23.1 (b)	20	20 (c)					
Sm		9.09 (b)	8.24	8.62 (c)				8.44 (c)	
Eu		1.77 (b)	1.64	1.74 (c)				1.33 (c)	
Gd		13.4 (b)							
Tb			2.11	2.18 (c)				2.2 (c)	
Dy		15.5 (b)						12.6 (c)	
Ho									
Er		9.02 (b)							
Tm									
Yb		8.36 (b)	7.94	8.21 (c)	9.6 (d)			9.8 (c)	
Lu			1.12	1.12 (c)				1.08 (c)	
Hf			7.37	7.93 (c)				6.8 (c)	
Ta			1.4	1.4 (c)				1.32 (c)	
W ppb								120 (c)	
Re ppb									
Os ppb									
Ir ppb			< 5	5 (c)					
Pt ppb									
Au ppb			< 6	< 7 (c)					
Th ppm			1.1	0.5 (c)		0.91	0.87 (e)	0.61 (c)	
U ppm		0.21 (b)	0.23	0.3 (c)		0.248	0.22 (e)	0.22 (c)	

technique: (a) XRF, (b) IDMS, (c) INAA, (d) "microchemical", (e) radiation count.



### References for 75061

Becker R.H. and Clayton R.N. (1975) Nitrogen abundances and isotopic compositions in lunar samples. *Proc. 6<sup>th</sup> Lunar Sci. Conf.* 2131-2149.

Butler P. (1973) Lunar Sample Information Catalog Apollo 17. Lunar Receiving Laboratory. MSC 03211 Curator's Catalog. pp. 447.

Fireman E.L., D'Amico J. and DeFelice J. (1973) Radioactivities vs. depth in Apollo 16 and 17 soil. *Proc. 4<sup>th</sup> Lunar Sci. Conf.* 2131-2144.

Graf J.C. (1993) Lunar Soils Grain Size Catalog. NASA Reference Pub. 1265, March 1993

Heiken G.H. (1974) A catalog of lunar soils. JSC Curator

Heiken G.H. (1975) Petrology of lunar soils. *Rev. Geophys. Space Phys.* **13**, 567-587.

Heiken G.H. and McKay D.S. (1974) Petrology of Apollo 17 soils. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 843-860.

Keith J.E., Clark R.S. and Bennett L.J. (1974a) Determination of natural and cosmic ray induced radionuclides in Apollo 17 lunar samples. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 2121-2138.

Hintenberger H., Schultz L., and Weber H.W. (1975) A comparison of noble gases in lunar fines and soil breccias: Implications for the origin of soil breccias. *Proc. 6<sup>th</sup> Lunar Sci. Conf.* 2261-2270.

Hubner W., Kirsten T and Kiko J. (1975) Rare gases in Apollo 17 soils with emphasis on analysis of size and mineral fractions of soil 74241. *Proc. 6<sup>th</sup> Lunar Sci. Conf.* 2009-2026.

Keith J.E., Clark R.S. and Bennett L.J. (1974a) Determination of natural and cosmic ray induced radionuclides in Apollo 17 lunar samples. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 2121-2138.

Korotev R.L. and Kremser D. (1992) Compositional variations in Apollo 17 soils and their relationships to the geology of the Taurus-Littrow site. *Proc. 22<sup>nd</sup> Lunar Planet. Sci. Conf.* 275-301.

LSPET (1973a) Apollo 17 lunar samples : Chemical and petrographic description. *Science* **182**, 659-690.

LSPET (1973c) Preliminary examination of lunar samples. Apollo 17 Preliminary Science Report. NASA SP-330, 7-1—7-46.

McKay D.S., Fruland R.M. and Heiken G.H. (1974) Grain size and the evolution of lunar soils. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 887-906.

Meyer C. (1973) Apollo 17 Coarse Fines (4-10 mm) Sample Location, Classification and Photo Index. Curator Report. pp. 182.

Mitchell J.K., Carrier W.D., Costes N.C., Houston W.N., Scott R.F. and Hovland H.J. (1973) 8. Soil-Mechanics. In Apollo 17 Preliminary Science Rpt. NASA SP-330. pages 8-1-22.

Morris R.V. (1976) Surface exposure indices of lunar soils: A comparative FMR study. *Proc. 7<sup>th</sup> Lunar Sci. Conf.* 315-335.

Morris R.V., Score R., Dardano C. and Heiken G. (1983) Handbook of Lunar Soils. Two Parts. JSC 19069. Curator's Office, Houston

Morris R.V. (1978) The surface exposure (maturity) of lunar soils: Some concepts and Is/FeO compilation. *Proc. 9<sup>th</sup> Lunar Sci. Conf.* 2287-2297.

- Müller O. (1974a) Solar wind nitrogen and indigenous nitrogen in Apollo 17 lunar samples. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 1907-1918.
- Müller O. (1974b) Solar wind and indigenous nitrogen in Apollo 17 lunar samples (abs). *Lunar Sci.* **V**, 534-536. Lunar Planetary Institute, Houston.
- Müller O. (1975) Lithophile trace and major elements in Apollo 16 and 17 lunar samples. *Proc. 6<sup>th</sup> Lunar Sci. Conf.* 1303-1312.
- Norris S.J., Swart P.K., Wright I.P., Grady M.M. and Pillinger C.T. (1983) A search for a correlatable, isotopically light carbon and nitrogen components in lunar soils and breccias. *Proc. 14<sup>th</sup> Lunar Planet. Sci. Conf.* in *J. Geophys. Res.* **88**, B200-B210.
- Nyquist L.E., Bansal B.M., Wiesmann H. and Jahn B.M. (1974) Taurus-Littrow chronology: Some constraints on the Early Lunar crustal development. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 1515-1540.
- Papike J.J., Simon S.B. and Laul J.C. (1982) The lunar regolith: Chemistry, Mineralogy and Petrology. *Rev. Geophys. Space Phys.* **20**, 761-826.
- Petrowski C., Kerridge J.F. and Kaplan I.R. (1974) Light element geochemistry of the Apollo 17 site. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 1939-1948.
- Rancitelli L.A., Perkins R.W., Felix W.D. and Wogman N.A. (1974a) Solar flare and lunar surface process characterization at the Apollo 17 site. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 2185-2203.
- Rhodes J.M., Rodgers K.V., Shih C., Bansal B.M., Nyquist L.E., Wiesmann H. and Hubbard N.J. (1974) The relationships between geology and soil chemistry at the Apollo 17 landing site. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 1097-1117.
- Rose H.J., Cuttitta F., Berman S., Brown F.W., Carron M.K., Christian R.P., Dwornik E.J. and Greenland L.P. (1974a) Chemical composition of rocks and soils at Taurus-Littrow. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 1119-1133.
- Schonfeld E. (1974) The contamination of lunar highland rocks by KREEP: Interpretations by mixing models. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 1269-1286.
- Wiesmann H. and Hubbard N.J. (1975) A compilation of the Lunar Sample Data Generated by the Gast, Nyquist and Hubbard Lunar Sample PI-Ships. Unpublished. JSC
- Wolfe E.W., Bailey N.G., Lucchitta B.K., Muehlberger W.R., Scott D.H., Sutton R.L. and Wilshire H.G. (1981) The geologic investigation of the Taurus-Littrow Valley: Apollo 17 Landing Site. US Geol. Survey Prof. Paper, 1080, pp. 280.